

Enzymatic and non-enzymatic oxidants and antioxidants involved in defense mechanisms against root-knot, reniform and citrus nematodes in their hosts

Afifi, A.M.R.*; Al-Sayed, A.A.**; Mahfoud, N.A.M² and Farahat, A.A.**

*Department of Biochemistry, Faculty of Agriculture- Cairo University

**Department of Zoology and Agricultural Nematology, Faculty of Agriculture- Cairo University

Abstract

A comparison between the changes in oxidants and antioxidants (involved in defense mechanisms in plants against pathogens) in cowpea, eggplant, jasmine, papaya and sour orange in response to infection with *Meloidogyne incognita*, *Rotylenchulus reniformis* and *Tylenchulus semipenetrans* was held. Different hosts react the same way to nematode infection, whereas oxidants, lipid peroxidase (MDA) and hydrogen peroxide (H₂O₂) were increased after nematode infection. Also antioxidant substances, Glutathione (GSH) and ascorbic acid (AA) as well as enzymes, superoxide dismutase (SOD), Catalase (CAT), ascorbate peroxidase (APX) and glutathione-S-transferase (GST) were also increased. The rates of increase differed according to nematode species, host plant and nematode initial population.

Key words: Oxidants, antioxidant enzymes, *Meloidogyne incognita*, *Rotylenchulus reniformis*, *Tylenchulus semipenetrans*.

Introduction

The root-knot nematodes *Meloidogyne* spp are the most important nematode pests worldwide due to their great damage resulted on the very wide host range which include more than 2000 plant species (**Gugino et al., 2008**). The reniform nematode, *Rotylenchulus reniformis* came after the root-knot nematode especially in the warmer parts of the world causing great losses in different field crops, vegetables, ornamentals and some orchards of fruit trees. The citrus nematode, *Tylenchulus semipenetrans*, the nematode pest responsible for the slow decline in citrus orchards and some other fruit trees like grape and olive all over the world. All the three nematode genera depend upon their development on their hosts on the formation of feeding sites eg. Giant cells, syncytia, hypertrophied cells and nurse cells. The success of nematode reproduction on their compatible hosts depend on the successful formation of such feeding sites which rely on the availability of certain concentrations of some chemicals and enzymes (**Baldacci-cresp et al., 2012**) to be available in host tissues.

In plants attacked by nematodes selective changes occur in the metabolism either as consequence of the establishment of a susceptible host-pathogen

interaction or as a result of resistance between host and parasite. Several models for resistance/susceptibility have been developed based on biochemical changes (**Kaplan and Keen, 1980; Giebel, 1974 and Zacheo et al., 1987**). There are many reports of enhanced peroxidases, polyphenoloxidase, and ascorbic acid oxidase following the interaction of nematodes with their hosts especially the resistant ones and this has led to the hypothesis that these enzymes may be important in the defense mechanism of host (**Brenneman and Black, 1979, Lazarovits and Ward, 1982; Zacheo et al., 1983, Saeed, 2005; Siahpoush et al., 2011, Aryal et al., 2011 and El-Belatagi et al, 2012**).

Generally, incompatibility to nematodes expressed after infection and active mechanisms involved compounds produced postinfectiously rather than performed constitutive plant products (**Kaplan and Keen, 1980**). Accordingly, plants develop defense mechanisms right away after nematode invasion. Most of these defense mechanisms are incompatible resistant interactions between plants and pathogens of which the formation of reactive oxygen species (ROS) are common (**Montes et al., 2004 and Bakker et al., 2006**). Such reactive oxygen species induced lipid peroxidation accounting for cell death after pathogen invasion. H_2O_2 acts as signaling molecule that triggers gene activation, or as cofactor in a process that requires new gene expression for both localized cell death and induction of defense genes in adjacent cells (**Mellersh et al., 2002**) as well as its direct effect on nematode development (**Karajeh, 2008 & Siahpoush, 2011**). Infected plants exhibit both enzymatic and non enzymatic antioxidant defense systems to frustrate ROS upon nematode infection. The accumulation of such materials in root tissues enhanced resistance in plants against invasion with new nematode larvae (**El-Belatagi et al., 2012**), of these Antioxidants GSH, SOD, Catalase and Ascorbate oxidase.

The present study is carried out to compare the changes in enzymatic and non enzymatic oxidants and antioxidants involved in defense mechanisms of cowpea, eggplant, jasmine, papaya and sour orange against the root-knot, the reniform and the citrus nematodes.

Materials and Methods

Stock cultures:

Pure cultures of the roo-knot nematode, *Meloidogyne incognita* (**Chitwood, 1949**), the reniform nematode, *Rotylenchulus reniformis* (**Linford and Olievera, 1940**) were obtained from the stock cultures belong to Nematology Research Center, Zoology and Agric. Nematology Dept. Faculty of Agric. Cairo Univ. These cultures are maintained on eggplant and pigeon pea, respectively. Pure culture of the citrus nematode, *Tylenchulus semipenetrans* (**Cobb, 1913**) was obtained from a citrus orchard at the Farm of the Fac. of Agric. Cairo Univ., Giza Egypt.

Test plants:

Cowpea (*Vigna sinensis*) cv Kareem 7, eggplant (*Solanum melongena*) hyb. Oneta F1, Papaya (*Carica papaya*) cv Solo, Jasmine (*Jasminum grandiflora*) cv Balady and sour orange (*Citrus aurantium*) were used in the present study.

One single seedling of each host crop grown in 15 cm diameter clay pots filled with sandy loam soil (1:1 v/v) were inoculated with 2000 or 4000 infective stage of one of the tested nematode species. Treatments were replicated 8 times as well as another untreated 8 to serve as a check. Pots of all treatments were arranged on a clean bench in a greenhouse at $30\pm 5^{\circ}\text{C}$ for 6 weeks and horticulturally treated the same.

Nematode extraction:

At the end of the experimental time, 4 pots of each treatment were soaked lonely in a plastic bucket half full of tap water until the root system could be separated. Each root system was stored in 5% formaldehyde in plastic jars. Soil suspension was quite stirred and nematodes were extracted using sieves and Baermann technique. Soil population was calculated using Hawkesly counting slide. Root population was determined under stereo binocular after staining the root with acid fuchsine (**Goody, 1957**). Fresh and dry samples of the inoculated and non inoculated plants of the other 4 replicates were sent to chemistry Laboratory, Department of Biochemistry, Fac. Of Agric. Cairo Univ. for determination of oxidant and antioxidant substances and enzymes.

Chemical analysis:**1. Determination of oxidative burst:****a. Lipid peroxidation (MDA contents)**

Thiobarbituric acid reaction (TBA) as described by **Heath and Parker (1968)**. The MDA equivalent was derived from the absorbance according to **Hodges et al. (1999)**.

b. Assay of hydrogen peroxide concentration:

Hydrogen peroxide was measured by the method described by Capaldi and Taylor (1983), with slight modification. The ground samples in 5% TCA (2.5 ml per 0.5 g fresh shoots or roots) with 50 mg active charcoal at 0°C , and centrifugated for 10 min. at $15,000 \times g$. Supernatant was collected, neutralized with 4 N KOH to pH 3.6 and used for H_2O_2 assay. The reaction mixture contained 200 μl of leaf extract, 100 μl of 3.4 mM 3- methylbenzothiazoline hydrazone (MBTH). The reaction was inhibited by adding 500 μl of horseradish peroxidase solution (90 μ per 100 ml) in 0:2 M sodium acetate (pH 3.6). Two minutes later 1400 μl of 1N HCL was added. Absorbance was read at 630 nm after 15 min.

2. Determination of antioxidant substances:

a. Determination of total glutathione:

The level of total acid-soluble SH compound (Glutathione GSH) was determined with Ellman's reagent according to **De Vos et al. (1992)**.

b. Ascorbic acid determination:

Levels of AA were determined following the procedure described by **Singh et al. (2006)** with few modifications. Fresh shoots or roots sample of a known weight (1g) was extracted with 3 ml of 5% (w/v) trichloroacetic acid (TCA) and centrifuged at 18,000 x g for 15 min. AsA was determined in a reaction mixture consisting of 0.2 ml of supernatant, 0.5 ml of 150 mM phosphate buffer (pH 7.4, containing 5 mM EDTA) and 0.2 ml of deionized water. Color was developed in reaction mixtures with the addition of 0.4 ml of 10% (w/v) TCA, 0.4 ml of 44% (v/v) phosphoric acid, 0.4 ml of α - α -dipyridyl in 70 % (v/v) ethanol and 0.2 ml of 3% (w/v) FeCl_3 . The reaction mixtures were incubated at 40°C for 40 min and the absorbance was read at 532 nm.

3. Determination of antioxidant enzymes:

- a. **Assay of SOD activity (SOD; EC.1.15.1.1):** The activity of SOD was assayed by measuring its ability to inhibit the photochemical reduction of NBT using the method of **Beauchamp and Fridovich (1971)**.
- b. **Assay of ascorbic peroxidase APX activity (APOX; EC.1.11.1.11):** Ascorbate peroxidase activity was estimated according to the method of **Nakano and Asada (1981)**.
- c. **Assay of catalase activity (CAT; EC 1.11.1.6):** Catalase activity was determined by consumption of H_2O_2 using the method of **Dhindsa et al. (1981)**.
- d. **Assay of Glutathione-S-transferase (GST; EC. 2.5.1.18):** Glutathione-S-transferase activity was measured according to the method of **Mannervik and Guthenberg (1981)**.

Results

The reproduction pattern and the rate of build up of the root-knot, *M. incognita*; the reniform, *R. reniformis* and the citrus nematode, *T. semipenetrans* on eggplant, papaya, cowpea, jasmine and sour orange as initially inoculated with 2000 or 4000 infective stage/plant are illustrated in Table (1). Data showed significant variations in the number of galls, egg-masses and final populations between the two initial inoculum levels of nematodes. The rate of nematode build up decreased in all cases by increasing the initial inoculums from 2000 to 4000 infective stage/pot. The highest rate of build up and the significant highest final

populations of *M. incognita* were achieved on eggplant and papaya, while those of *R. reniformis* were achieved on jasmine and cowpea. These results evinced that eggplant and papaya were preferable hosts to *M. incognita* than jasmine and cowpea which were more preferable to *R. reniformis*. The citrus nematode reproduced normally on sour orange.

Table (1): Infectivity and build-up of the root-knot, the reniform nematode and the citrus nematode on different host plants

Host	Inoculum level	Counts of <i>Meloidogyne incognita</i>		
		Galls	Final population	Pf/Pi
Eggplant	2000	1146 b	7284 c	3.6
	4000	1244 a	10488 a	2.6
Papaya	2000	896 d	5648 d	2.8
	4000	1022 c	7793 b	1.9
Cowpea	2000	311 f	4250 f	2.1
	4000	451 e	5500 e	1.4
Jasmine	2000	469 e	2926 h	1.5
	4000	522 e	3763 g	0.9
Counts of <i>Rotylenchulus reniformis</i>				
		Egg-masses	Final population	Pf/Pi
Eggplant	2000	207 h	4380 h	2.2
	4000	243 g	6398 f	1.6
Papaya	2000	274 f	6321 g	3.2
	4000	402 e	8164 e	2.0
Cowpea	2000	1013 d	9843 d	4.9
	4000	1689 b	13066 c	3.3
Jasmine	2000	1547 c	16044 b	8.0
	4000	1969 a	20659 a	5.1
Counts of <i>Tylenchulus semipenetrans</i>				
		Egg-masses	Final population	Pf/Pi
Sour orange	2000	255	3205	1.6
	4000	292 **	3594 **	0.9

Concerning the changes in plant enzymes involved in defense mechanisms against plant pathogens, data in table (2) indicated that lipid peroxidase (MDA) and hydrogen peroxide (H₂O₂) in shoots and roots of healthy cowpea, eggplant, jasmine,

Table (2): Comparison between the contents of shoots and roots of different host plants of oxidants, lipid peroxidation (MDA) and hydrogen peroxide (H₂O₂) in response to infection with different nematode species and inoculum levels.

Host	Nematode	Inoc. Level	MDA $\mu\text{mol/g fw}$				H ₂ O ₂ $\mu\text{mol/g fw}$			
			Shoot	% inc.	Root	% inc.	Shoot	% inc.	Root	% inc.
Cowpea	Control		2.50 ^{pp} ±0.36		3.08 ^{pp} ±0.07		4.06 ^m ±0.08		5.86 ^m ±0.07	
	<i>M. incog.</i>	2000	2.56 ^{pp} ±0.14	2.4	3.32 ^{mm} ±0.13	7.8	5.75 ^r ±0.10	41.6	7.48 ^{mm} ±0.13	26.6
	<i>M. incog.</i>	4000	6.05 ^r ±0.03	142.0	7.55 ^r ±0.27	145.1	13.55 ^r ±0.17	233.7	19.41 ^r ±0.14	231.2
	<i>R. renifo.</i>	2000	3.29 ^{mm} ±0.07	31.6	3.95 ^{rk} ±0.02	28.3	8.50 ^o ±0.17	109.4	9.83 ^{rk} ±0.17	64.3
Eggplant	<i>R. renifo.</i>	4000	7.92 ^r ±0.08	216.8	8.50 ^o ±0.36	176.0	17.99 ^r ±0.08	343.1	24.40 ^o ±0.12	316.4
	Control		2.44 ^r ±0.04		2.92 ^o ±0.05		4.54 ^o ±0.10		6.09 ^{mm} ±0.03	
	<i>M. incog.</i>	2000	6.41 ^h ±0.14	162.7	7.01 ^h ±0.08	140.1	16.54 ^h ±0.18	264.3	18.24 ^h ±0.10	199.5
	<i>M. incog.</i>	4000	12.85 ^h ±0.03	418.4	13.98 ^h ±0.06	378.8	38.89 ^h ±0.06	756.6	45.89 ^h ±0.07	653.5
Jasmine	<i>R. renifo.</i>	2000	2.83 ^{pp} ±0.09	18.0	2.96 ^o ±0.04	1.4	4.69 ^{mm} ±0.08	3.3	6.11 ^{mm} ±0.10	0.3
	<i>R. renifo.</i>	4000	5.59 ^h ±0.02	129.1	6.04 ^h ±0.08	106.9	9.80 ^h ±0.07	115.9	10.55 ^h ±0.07	73.2
	R+M	2000	2.87 ^{pp} ±0.07	17.6	3.03 ^o ±0.08	3.8	7.68 ^h ±0.08	69.2	9.94 ^h ±0.02	63.2
	R+M	4000	6.99 ^h ±0.03	186.5	8.06 ^h ±0.06	176.0	13.56 ^h ±0.07	198.7	16.40 ^h ±5.26	169.3
Papaya	Control		2.66 ^{pp} ±0.05		3.23 ^{mm} ±0.04		4.80 ^o ±0.06		6.13 ^{mm} ±0.10	
	<i>M. incog.</i>	2000	2.98 ^{pp} ±0.02	12.0	3.88 ^h ±0.08	20.1	8.46 ^h ±0.09	74.1	9.83 ^h ±0.10	51.1
	<i>M. incog.</i>	4000	6.63 ^h ±0.08	149.3	7.82 ^h ±0.08	142.1	18.40 ^h ±0.09	278.6	21.60 ^h ±0.12	252.9
	<i>R. renifo.</i>	2000	4.89 ^h ±0.07	83.8	5.59 ^h ±0.06	73.1	11.08 ^h ±0.04	128.0	13.80 ^h ±0.07	121.9
Citrus	<i>R. renifo.</i>	4000	11.00 ^h ±0.26	313.5	12.05 ^h ±0.35	273.3	30.11 ^h ±0.01	519.6	35.56 ^h ±0.07	480.1
	R+M	2000	2.98 ^{pp} ±0.03	12.0	3.96 ^{rk} ±0.06	22.6	9.61 ^h ±0.11	97.7	9.94 ^h ±0.03	62.2
	R+M	4000	7.22 ^h ±0.07	171.4	8.53 ^h ±0.25	164.1	19.83 ^h ±0.13	308.0	23.10 ^h ±0.48	276.8
	Control		2.80 ^{pp} ±0.07		3.03 ^o ±0.08		4.36 ^h ±0.10		5.96 ^{mm} ±0.12	
Citrus	<i>M. incog.</i>	2000	3.95 ^h ±0.26	41.1	4.19 ^h ±0.23	38.3	10.91 ^h ±0.10	150.2	12.11 ^h ±0.20	103.2
	<i>M. incog.</i>	4000	8.13 ^h ±0.05	190.4	9.82 ^h ±0.02	224.1	23.20 ^h ±0.06	432.1	29.13 ^h ±0.01	388.8
	<i>R. renifo.</i>	2000	3.09 ^{mm} ±0.03	10.4	3.62 ^h ±0.16	19.5	4.96 ^h ±0.11	13.8	6.33 ^{mm} ±0.10	6.2
	<i>R. renifo.</i>	4000	6.69 ^h ±0.05	138.9	7.56 ^h ±0.13	149.5	10.13 ^h ±0.02	132.3	13.46 ^h ±0.08	125.8
Citrus	Control		3.01 ^{mm} ±0.17		3.90 ^h ±0.06		5.11 ^h ±0.09		6.36 ^{mm} ±0.09	
	<i>T. semi.</i>	2000	3.11 ^{mm} ±0.16	3.32	3.56 ^{mm} ±0.11	0.0	7.12 ^h ±0.11	39.3	8.28 ^h ±0.07	30.2
	<i>T. semi.</i>	4000	6.90 ^h ±0.07	129.2	7.59 ^h ±0.15	94.6	15.56 ^h ±0.08	204.4	21.60 ^h ±0.11	239.6

*Values followed by the same letter are not significantly different ($p > 0.5$)

papaya and sour orange were at the lowest levels and varied significantly in the majority of cases with those infected with *M. incognita*, *R. reniformis* or *T. semipenetrans*. Nematode infection increased levels of MDA and H₂O₂ in all plants. In all cases, higher inoculum levels (4000/ plant) resulted in significant higher increases in MDA and H₂O₂. The highest significant increase was observed in shoots and roots of eggplant infected with 4000 J₂ of *M. incognita*, followed by those in shoots and roots of jasmine infected with *R. reniformis* then in shoots and roots of papaya infected with *M. incognita* at the same inoculum level. Sour orange behaved the same in response to infection with *T. semipenetrans*. MDA and H₂O₂ increased significantly especially at the high inoculum level.

Non enzymatic antioxidants, glutathione (GSH) and ascorbic acid significantly increased in response to nematode infection. The highest significant increase was observed in shoots and roots of eggplant infected with 4000 infective stages of *R. reniformis* followed by those in shoots and roots of cowpea infected with 4000 J₂ of *M. incognita* then those in papaya infected with 4000 infective stage of *R. reniformis* (Table 3). It could be generally observed that the increase in such antioxidants depends on nematode species, inoculum level and host plant. Similarly, the citrus nematode resulted in increasing GSH and ascorbic acid significantly as compared to healthy plants.

With the activity of antioxidant enzymes, superoxide dismutase (SOD), ascorbate peroxidase (APX) and catalase (CAT), data in table 4 indicated that the activities of such enzymes in healthy plants were at low levels without significant differences between different hosts. Significant increase was observed as a result of nematode infection, *Rotylenchulus reniformis* on eggplant at 4000 infective stage achieved the highest activity of the three enzymes which varied significantly with other nematodes on other hosts. *Meloidogyne incognita* at the same inoculum level on cowpea came after, followed by *R. reniformis* on papaya, *M. incognita* on jasmine then *T. semipenetrans* on sour orange (Tables 4, 5).

Glutathione-S-transferase (GST) activity in shoots and roots of different hosts was likely increased significantly by nematode infection. *Meloidogyne incognita* at the high inoculum level recorded the highest significant activity in shoots and roots of eggplant, followed by *Rotylenchulus reniformis* in shoots and roots of jasmine and then shoots and roots of cowpea. GST activity was also increased significantly in shoots and roots of sour orange as a result of infection with *T. semipenetrans* (Table 5).

Infection of eggplant and jasmine with concomitant population of R+M resulted in higher rates of oxidants and antioxidants comparing to infection of eggplant with single population of *R. reniformis* or jasmine with single population of *M. incognita* at the same inoculum level.

Table (3): Comparison between the contents of shoots and roots of different host plants of antioxidant substances, glutathione (GSH) and ascorbic acid (AA) in response to infection with different nematode species and inoculum levels.

Host	Nematode	Inoc. Level	GSH $\mu\text{mol/g fw}$			Ascorbic acid mg/g fw				
			Shoot	Root	% inc.	Shoot	Root	% Inc.		
Cowpea	Control		15.10 ^a ±0.17		17.9 ^p ±0.10		8.55 ^q ±0.26		11.71 ^r ±0.17	
	<i>M. incog.</i>	2000	24.9 ^b ±0.20	64.9	29.7 ^q ±0.20	65.9	21.40 ^r ±0.36	150.3	25.30 ^s ±0.36	116.1
	<i>M. incog.</i>	4000	45.4 ^c ±0.62	200.7	55.8 ^r ±0.26	210.6	28.93 ^s ±0.20	238.4	35.77 ^t ±0.26	205.5
	<i>R. renifo.</i>	2000	21.1 ^b ±0.55	39.7	23.2 ^q ±0.45	29.6	14.50 ^r ±0.26	89.6	17.21 ^r ±0.26	47.0
	<i>R. renifo.</i>	4000	30.8 ^c ±0.28	102.7	35.96 ^r ±0.10	100.9	19.55 ^r ±0.26	128.7	25.14 ^r ±0.26	114.7
Eggplant	Control		14.4 ^a ±0.36		16.8 ^a ±0.26		8.95 ^a ±0.26		10.88 ^a ±0.26	
	<i>M. incog.</i>	2000	13.4 ^a ±0.28	0.00	16.1 ^a ±0.65	0.00	8.60 ^a ±0.17	0.0	10.20 ^a ±0.36	0.0
	<i>M. incog.</i>	4000	16.7 ^{ab} ±0.20	16.0	19.8 ^a ±0.26	17.9	11.68 ^a ±0.26	30.5	13.57 ^a ±0.20	24.7
	<i>R. renifo.</i>	2000	31.9 ^b ±0.26	121.5	36.5 ^a ±0.20	117.3	23.50 ^a ±0.17	162.6	28.30 ^b ±0.36	160.1
	<i>R. renifo.</i>	4000	56.7 ^c ±0.28	293.8	68.8 ^a ±0.26	309.5	32.92 ^a ±0.36	267.8	37.73 ^a ±0.20	246.8
Jasmine	Control		14.1 ^a ±0.17		15.9 ^a ±0.10		9.23 ^a ±0.36		11.51 ^a ±0.20	
	<i>M. incog.</i>	2000	19.7 ^a ±0.20	39.7	21.9 ^{ab} ±0.26	37.7	18.60 ^b ±0.17	101.5	21.60 ^b ±0.20	87.7
	<i>M. incog.</i>	4000	32.1 ^b ±0.36	127.7	43.6 ^a ±0.20	174.2	22.93 ^b ±0.26	148.4	30.77 ^b ±0.26	167.3
	<i>R. renifo.</i>	2000	14.0 ^a ±0.52	0.0	16.9 ^a ±0.26	6.3	11.30 ^{ab} ±0.26	22.4	12.40 ^a ±0.36	7.7
	<i>R. renifo.</i>	4000	18.2 ^a ±0.17	29.1	22.4 ^a ±0.45	40.9	13.90 ^{ab} ±0.20	50.6	15.80 ^a ±0.26	37.3
Papaya	Control		13.9 ^a ±0.17		16.2 ^a ±0.36		8.37 ^a ±0.36		10.92 ^a ±0.26	
	<i>M. incog.</i>	2000	16.1 ^a ±0.65	15.8	18.2 ^a ±0.36	12.3	11.60 ^a ±0.17	38.6	16.70 ^{ab} ±0.20	52.9
	<i>M. incog.</i>	4000	21.1 ^b ±0.36	51.8	24.4 ^a ±0.26	50.6	19.33 ^b ±0.36	130.9	23.24 ^b ±0.55	112.8
	<i>R. renifo.</i>	2000	26.3 ^b ±0.55	89.2	32.5 ^a ±0.26	100.6	21.50 ^a ±0.17	156.9	25.20 ^a ±0.36	130.8
	<i>R. renifo.</i>	4000	41.4 ^c ±0.26	197.8	47.3 ^a ±0.43	192.0	27.51 ^a ±0.26	228.7	33.27 ^a ±0.40	204.7
Citrus	Control		16.22 ^{ab} ±0.10		18.64 ^a ±0.11		9.32 ^a ±0.13		11.73 ^a ±0.09	
	<i>T. semi.</i>	2000	21.3 ^b ±0.55	31.3	24.3 ^a ±0.55	30.4	15.20 ^a ±0.36	63.09	19.80 ^a ±0.26	68.8
	<i>T. semi.</i>	4000	35.9 ^c ±0.08	121.3	38.73 ^a ±0.09	54.1	24.28 ^a ±0.07	160.5	29.59 ^a ±0.02	155.3

*Values followed by the same letter are not significantly different ($p=0.5$)

Table (4): Comparison between the activities of antioxidant enzymes, Super oxide dismutase (SOD) and ascorbate peroxidase (APX) in shoots and roots of different host plants in response to infection with different nematode species and inoculum levels.

Host	Nematode	Inoc. level	SOD (Unit/g fw)				APX (Unit/mg fw)			
			Shoot	% Inc.	Root	% Inc.	Shoot	% Inc.	Root	% Inc.
Cowpea	Control		89.1 ^m ±1.05		73.3 ^a ±0.45		14.9 ^a ±0.26		11.3 ^a ±0.36	
	<i>M. incog.</i>	2000	136.9 ^{op} ±0.26	53.7	119.9 ^f ±0.26	63.6	24.6 ^b ±0.26	65.1	19.4 ^b ±0.36	71.7
	<i>M. incog.</i>	4000	186.8 ^{op} ±0.26	109.7	153.4 ^e ±0.26	109.3	35.4 ^b ±0.36	137.6	31.7 ^b ±0.20	180.5
	<i>R. renifo.</i>	2000	112.6 ^{pk} ±0.26	26.4	98.6 ^g ±0.26	34.5	19.8 ^{mm} ±0.26	32.9	14.4 ^m ±0.26	27.4
	<i>R. renifo.</i>	4000	159.3 ^{opq} ±0.36	78.8	143.1 ^f ±0.36	95.2	26.9 ^a ±0.26	80.5	22.5 ^f ±0.20	99.1
Eggplant	Control		93.1 ^{kim} ±0.36		84.3 ^q ±0.36		16.1 ^a ±0.65		12.9 ^{op} ±0.26	
	<i>M. incog.</i>	2000	103.8 ^{kj} ±0.41	11.50	86.3 ^q ±0.36	14.8	16.7 ^a ±0.26	3.7	13.2 ^{osp} ±0.60	2.3
	<i>M. incog.</i>	4000	111.8 ^{ks} ±1.99	20.0	100.3 ^m ±0.26	19.0	17.9 ^b ±0.20	11.2	13.8 ^{mm} ±0.26	7.0
	<i>R. renifo.</i>	2000	143.2 ^{opq} ±0.45	53.8	129.1 ^f ±0.65	53.1	28.3 ^f ±0.26	75.8	25.1 ^a ±0.65	94.6
	<i>R. renifo.</i>	4000	202.5 ^a ±0.43	117.5	196.7 ^a ±0.35	133.2	41.0 ^a ±0.26	154.7	38.5 ^a ±0.20	198.5
	R+M	2000	117.9 ⁿⁱ ±0.20	26.6	108.6 ^f ±0.26	28.8	20.5 ^a ±0.36	27.3	17.0 ^a ±0.75	31.8
	R+M	4000	174.5 ^{bc} ±0.20	87.4	139.6 ^b ±0.45	65.6	28.5 ^a ±0.36	77.0	21.3 ^a ±0.55	65.1
Jasmine	Control		88.3 ^m ±0.26		71.5 ^f ±0.20		15.7 ^a ±0.26		12.6 ^p ±0.20	
	<i>M. incog.</i>	2000	129.9 ^{opm} ±0.26	46.6	110.3 ^f ±0.41	54.3	20.0 ^a ±0.45	27.4	16.1 ^a ±0.55	27.8
	<i>M. incog.</i>	4000	173.3 ^{bc} ±0.55	95.6	159.0 ^e ±2.64	122.4	29.5 ^a ±0.26	87.9	25.7 ^a ±0.36	104.0
	<i>R. renifo.</i>	2000	93.6 ^{mm} ±0.20	5.6	84.3 ^q ±0.36	17.9	15.3 ^f ±0.36	0.0	11.5 ^a ±0.36	0.0
	<i>R. renifo.</i>	4000	116.1 ^f ±0.65	31.0	108.8 ^f ±0.10	52.2	19.4 ^m ±0.26	23.6	15.5 ^b ±0.36	23.0
	R+M	2000	112.8 ^{ks} ±0.26	27.3	96.1 ^g ±0.65	34.4	17.6 ^{op} ±0.26	12.1	13.8 ^{mm} ±0.36	9.5
	R+M	4000	156.1 ^{op} ±0.52	76.2	141.6 ^a ±0.26	98.0	22.5 ^a ±0.26	43.3	18.3 ^a ±0.36	45.2
Papaya	Control		90.7 ^{km} ±0.20		82.1 ^f ±0.36		15.4 ^a ±0.36		12.8 ^p ±0.26	
	<i>M. incog.</i>	2000	103.6 ^{kj} ±0.20	14.2	94.3 ^p ±0.36	14.9	17.3 ^a ±0.36	12.3	13.1 ^{osp} ±0.55	2.3
	<i>M. incog.</i>	4000	156.5 ^{op} ±0.26	72.6	139.9 ^f ±0.26	70.4	21.5 ^f ±0.20	39.6	15.2 ^a ±0.55	18.8
	<i>R. renifo.</i>	2000	132.6 ^{opm} ±0.26	46.2	119.7 ^f ±0.20	45.8	22.7 ^f ±0.20	47.4	18.9 ^{opm} ±0.26	47.7
	<i>R. renifo.</i>	4000	179.3 ^{bc} ±0.26	97.7	163.1 ^f ±0.36	98.7	33.8 ^f ±0.36	119.5	27.4 ^a ±0.26	114.1
Citrus	Control		92.9 ^{kmi} ±0.17		70.5 ^f ±0.26		15.4 ^a ±0.36		13.6 ^{ns} ±0.26	
	<i>T. sem i.</i>	2000	122.3 ⁿⁱ ±0.26	31.6	109.2 ^f ±0.55	54.9	22.4 ^a ±0.36	45.5	16.8 ^f ±0.17	23.5
	<i>T. sem i.</i>	4000	165.1 ^{op} ±0.75	77.7	150.7 ^a ±0.17	113.8	30.1 ^a ±0.55	95.5	26.3 ^a ±0.26	93.4

* Values followed by the same letter are not significantly different (p=0.5)


Table (5): Comparison between the activities of antioxidant enzymes, Catalase (CAT) and glutathione-S-transferase (GST) in shoots and roots of different host plants in response to infection with different nematode species and inoculum levels.

Host	Nematode	Inoc. level	CAT unit/g fw				GST unit/g fw			
			Shoot	% Inc.	Root	% Inc.	Shoot	% Inc.	Root	% Inc.
Cowpea	Control		34.4 w		24.6 v		5.56 v		6.11 y	
	<i>M. incog.</i>	2000	81.2 j	77.9	48.0 l	93.1	14.40 s	159.0	17.13 t	180.5
	<i>M. incog.</i>	4000	103.4 b	200.6	94.1 b	282.5	29.84 h	436.7	36.86 h	503.3
	<i>R. renifo.</i>	2000	47.1 o	36.9	38.9 o	58.1	18.33 o	229.7	20.88 o	238.5
	<i>R. renifo.</i>	4000	78.5 f	128.2	66.4 f	169.9	37.76 c	579.1	40.58 d	564.2
Eggplant	Control		39.8 t		29.3 t		5.16 w		6.36 x	
	<i>M. incog.</i>	2000	45.9 p	15.3	33.4 r	14.0	24.68 k	378.3	27.38 l	330.5
	<i>M. incog.</i>	4000	48.9 n	22.6	38.8 o	32.4	51.31 a	894.4	56.67 a	791.0
	<i>R. renifo.</i>	2000	68.2 i	71.6	54.8 j	87.0	12.66 t	145.4	15.84 u	149.1
	<i>R. renifo.</i>	4000	117.8 a	196.0	98.8 a	237.2	25.39 j	392.1	31.58 k	396.5
	R+M	2000	44.8 q	12.1	36.4 pq	24.2	16.25 r	214.9	19.13 s	200.8
	R+M	4000	78.6 f	97.5	64.7 g	120.8	28.40 i	450.4	36.32 i	471.1
Jasmine	Control		36.9 v		26.1 u		4.58 x		6.39 x	
	<i>M. incog.</i>	2000	53.9 l	46.1	44.4 m	70.1	17.31 p	278.0	19.40 r	203.6
	<i>M. incog.</i>	4000	90.3 d	144.7	78.5 d	200.8	31.51 e	588.0	38.65 f	504.9
	<i>R. renifo.</i>	2000	43.8 r	18.2	30.8 s	18.0	22.27 l	386.2	24.58 m	284.7
	<i>R. renifo.</i>	4000	56.09 k	52.0	43.6 n	67.1	43.51 b	850.0	49.57 b	675.7
	R+M	2000	45.0 q	22.0	36.1 q	38.3	18.76 n	309.6	20.35 p	218.5
Papaya	R+M	4000	70.5 h	91.1	61.1 i	134.1	33.37 d	628.6	39.48 e	517.9
	Control		42.1 s		30.7 s		4.50 x		6.95 w	
	<i>M. incog.</i>	2000	45.7 p	8.6	36.9 p	20.2	21.31 m	373.6	23.35 n	236.0
	<i>M. incog.</i>	4000	74.1 g	76.0	62.6 h	103.9	43.65 b	870.0	48.37 c	596.0
	<i>R. renifo.</i>	2000	61.0 j	44.9	48.9 k	59.3	16.82 q	273.8	19.11 s	175.0
Citrus	<i>R. renifo.</i>	4000	92.8 c	120.4	84.1 c	173.9	30.68 f	581.8	36.07 j	419.0
	Control		38.6 u		29.5 t		7.60 u		8.36 v	
	<i>T. semi.</i>	2000	50.9 m	31.8	43.6 n	47.8	16.26 r	113.9	19.84 q	137.3
	<i>T. semi.</i>	4000	84.5 e	118.9	73.3 e	148.5	30.43 g	300.4	38.25 g	357.5

*Values followed by the same letter are not significantly different ($p > 0.5$)

Comparing the percentage increase in oxidants, antioxidant substances and enzymes in perennial hosts in response to infection with 4000 infective stage of the three nematode species, data in Fig. (1) illustrate that the highest percentages of increase in peroxidase (MDA) and H₂O₂ were resulted in jasmine shoots and roots in response *R. reniformis* followed by those accomplished by *M. incognita* in papaya shoots and roots; however, the lowest rates of increase in such oxidants were recorded in sour orange shoots and roots infected with *T. semipenetrans*. Regarding the oxidant substances Glutathione (GSH) and ascorbic acid (AA) and enzymes SOD, APX and CAT the highest percentages of increase were observed in sour orange shoots and roots, followed by papaya and then jasmine in response to infection with *T. semipenetrans*, *R. reniformis* and *M. incognita*, respectively. In case of antioxidant enzymes, percentage increase in GST was the highest comparing to other antioxidant enzymes with superiority of *R. reniformis*, then *M. incognita* and *T. semipenetrans*.

Fig (1): % increase in oxidants and antioxidants in shoots and roots of jasmine, papaya and sour Orange in response to infection with, *R. reniformis*, *M. incognita* and *T. smipenetrans*, respectively.



It could be generally concluded that host plants respond to nematode infection, to great extent, by the same way. The oxidant and antioxidant enzymatic

and non enzymatic substances are involved in the defense mechanisms exerted by plants against different nematode pests. Yet, the amounts produced differed from one host to another, nematode species to another, host compatibility as well as the size of nematode population.

Discussion

It is generally known that incompatibility to nematodes expressed after infection and active mechanisms involved compounds produced postinfectiously rather than performed constitutive plant products (**Kaplan and Keen, 1980**). From this point of view, plants develop defense mechanisms right away after nematode invasion. Most of these defense mechanisms are incompatible resistant interactions between plants and pathogens of which the formation of reactive oxygen species (ROS) are common (**Montes *et al.*, 2004 and Bakker, *et al.*, 2006**). Such reactive oxygen species induced lipid peroxidation accounting for cell death after pathogen invasion. Hence, increasing the rates of MDA and H₂O₂ in different hosts in response to infection with *M. incognita*, *R. reniformis* and *T. semipenetrans* in the present study as compared to healthy plants accounted for the defense mechanism against nematode invasion. The rates of increase in both oxidants depend on nematode species and inoculum levels as well as host plant. Our results agreed with those of **Davis *et al.*, 2000 and Huang *et al.*, 2004** who said that the initial reaction of the susceptible cultivars is similar to that of resistant host and may be result from nematode secretions into plant tissues.

Infected plants exhibited both enzymatic and non-enzymatic antioxidant defense systems to frustrate ROS upon nematode infection. The significant increase of non-enzymatic antioxidants such as glutathione (GSH) and total ascorbic acid may be resulted from the enhancement of MDA and H₂O₂ production after nematode invasion. In the present study, infection with the three nematode species resulted in significant increase in GSH and ascorbic acid in the different host plants. The highest significant increase was observed in shoots and roots of eggplant infected with 4000 infective stages of *R. reniformis* followed by those in shoots and roots of cowpea infected with 4000 J₂ of *M. incognita*. The significant lower levels of both antioxidants in preferable hosts to both nematodes may be attributed to its consumption in scavenging higher levels of peroxidases produced in such hosts. The involvement of GSH in cellular metabolism and its potential agrobiotechnological application has been realized to resist biotic and abiotic stress and pests. Regulation of GSH considered to enhance pathogen resistance in plants and maintain antioxidant capacity of these substances in scavenging free radicals (**Afifi *et al.*, 2011; Hari *et al.*, 2011 and Afifi *et al.*, 2012**). The role of ascorbic acid in plant defense mechanisms against nematodes was illustrated by **Arrigoni *et al.* (1979)**. They reported that ascorbic acid depletion in plants attenuated resistance in tissues to nematode infections. They hypothesized that plants utilized ascorbic acid for synthesis of mitochondrial hydroxyproline proteins which control the development

of cyanide resistant respiration. They also stated that ascorbic acids synthesis was stimulated in resistant plants. **Arrigoni (1979)** also hypothesized that cyanide-resistant respiration (CCR), commonly associated with wounds is requisite to activations of biological defense mechanisms. Ascorbic acid-dependent synthesis of hydroxyproline-containing protein is associated with CCR in mitochondria. Cyanide resistant respiration generates hydrogen peroxide from which superoxide is generated by peroxidase. Superoxides are extremely toxic and diffuse through cells to oxidize functional groups of enzymes and phospholipids, reduce S-S bonds and cause macromolecule and membrane injuries.

Increase in superoxide dismutase (SOD) and peroxidase activity results to be an adaptive response which provides the plant with protection against biotic and abiotic stress (**Guida et al., 1992**). The protective activity of SOD, and catalase (CAT) was enhanced in susceptible but decreased in resistant plants (**Zacheo et al., 1993**). Superoxide dismutase prevents the deleterious effect of O₂ radicals in root cells and transform it to H₂O₂ which is then transformed by catalase to harmless O₂ + H₂O. Accordingly, in susceptible tomato roots infested with *M. incognita* SOD activity considerably increased in comparison to uninfected controls and decreased in resistant cultivars (**Zacheo et al., 1987**). These findings are in accordance with our results whereas superoxide dismutase (SOD), ascorbate oxidase (APX), Glutathione-S-transferase (GST) and catalase (CAT) increased after nematode infection. The increment rates varied according to host plant, nematode species and inoculum level.

References

- Afifiy, A.E.M.M.R.; El-Beltagi, H.S.; Abd El-Salam, Samiha.M. and Omran, Azza, A. (2012).** Biochemical changes in phenols, flavonoids, tannins, vitamin E, carotene and antioxidant activity during soaking of three white sorghum varieties. *Asian Pacific Journal of Tropical Biomedicine* (2012), 203-209.
- Afifiy, A.E.M.M.R.; El-Beltagi, H.S.; Fayed, S.A. and Shalaby, E.A. (2011).** Acridical activity of different extracts from *Syzygiumcumini* L. Seekls (Pomposia) against *Tetranychus urticae* Koch. *Asian Pacific Journal of Tropical Biomedicine* (2011), 359-364
- Arrigoni, O.; Zacheo, G.; Arrigoni-Liso, R.; Bleve-Zacheo, T. and Lamberti, F. (1979).** Relationship between ascorbic acid and resistance in tomato plants to *Meloidogyne incognita*. *Phytopathology*, 69: 570-581.
- Aryal, S.K., Davis, R.F., Stevenson, K.L., Timper, P. and Pingsheng, J. (2011).** *Meloidogyne incognita* enzymes involved in systemic acquired resistance. *J. Nematol.* 43(3-4): 152-159
- Bakker, E., Dees, R., Bakker, J., Goverse, A. (2006).** Mechanisms involved in plant resistance to nematodes. In Tuzen, S, Bent, E (eds) *Multigenic and Induced Systemic Resistance in Plants*. New York Springer Sci. pp. 314-344.

- Baldacci-Cresp, F.; Chang, C.; Macourt, M.; Deborde, C.; Hopkins, J.; Lecomte, P.; Bernillon, S.; Brouquisse, R.; Moing, A.; Abad, P.; Herouart, D.; Puppo, A.; Favery, B. and Frendo, P. (2012).** (Homo) glutathione deficiency impairs root-knot nematode development in *Medicago truncatula*. PLoS Pathogen / www. Plospathogen. Org. 8(1): e1002471.
- Beauchamp, C. and Fridovich, I. 1971.** Superoxide dismutase: improved assays and assay applicable to acrylamide gels. Anal. Biochem. 44: 276-287.
- Brenneman, J.A. and Black, L.L. 1979.** Respiration and terminal oxidases in tomato leaves infected by *Phytophthora infestans*. Physiol. Plant Pathol. 14:281-290.
- Capaldi, D.J. and Taylor, K.E. 1983.** A new peroxides color reaction: oxidative coupling of 3-methyl-2-benzothiazolinone hydrazone (MBTH) with its formaldehyde azine application to glucose and choline oxidases. Anal. Biochem., 129:329-336.
- Chitwood, B.G. (1949).** "Root-knot nematodes" Part 1. A revision of the genus *Meloidogyne* Goeldi, 1887. Proceeding of Helminthological Society of Washington, 16: 90-104.
- Cobb, C.A. (1913).** New nematode genera found inhabiting fresh water and non-brackish soils. Journal Wash. Acad., 3 : 432-444.
- Davis, E.L.; Hussey, R.S.; Baum, T.J.; Barker, J.; Schots, A.; Rosso, M.N. And Abad, P. (2000).** Nematode parasitism genes. Annual Review of Phytopathology, 38: 356-396.
- De Vos, C.H.; Vonk, M.; Vooijs, J.R. and Henk, S. 1992.** Glutathione depletion due to copper-induced phytochelatin synthesis causes oxidative stress in *Silene cucubalus*. Plant Physiology, 98: 859-858.
- Dhindsa, R.S.; Plumb-Dhindsa, P. and Thorpe, T.A. 1981.** Leaf senescence correlated with increased levels of membrane permeability and lipid peroxidation, and decrease levels of superoxide dismutase and catalase. J. Exp. Bot., 32: 93-101.
- Ei-Beltagi, H.S.; Farahat, A.A.; Al-Sayed, A.A. and Mahfoud, N.A.M. (2012).** Response of antioxidant substances and enzyme activities as defense mechanism against root-knot nematode infection. Not. Bot. Hort. Agrobot. Cluj 40(1): 132-142.
- Giebel, J. (1974).** Biochemical mechanisms of plant resistance to nematode: A review. Journal of Nematology, 6(4): 175-184.
- Goody, J.B. 1957.** Laboratory methods for work with plant soil nematodes. Bull. No.2. Ministry of Agriculture, London, pp. 47.
- Gugino, B.K.; Ludwig, J.W. and Abawi, G. S. 2008.** An on-farm bioassay for assessing *Meloidogyne hapla* infestations as a decision management tool. Crop Protection, 27: 785-791.
- Guida, G.; Zacheo, G and Bleve-Zacheo, T. (1992).** Activation of detoxifying enzymes in tomato roots following paraquat treatment and nematode

infection. *Nematol, medit.* (20): 203-209.

- Hari, C. M.; Gajbhiye, V.T. Singh, G. (2011).** Salicylic acid –induced glutathione status in tomato crop and resistance to root-knot nematode, *Meloidogyne incognita* (Kofoid & White) Chitwood. *J. Xenobiotics* 1(5): 22-28.
- Heath, R. and Packer, L. 1968.** Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. *Arch.Biochem. Biophys.*, 125:189-198.
- Hodges, D.M.; Delong, J.M.; Forney, C.F. and Prange, R.K. 1999.** Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta*, 207, 604-611.
- Huang, G.; Dong,R.; Maier,T.; Allen, R.; Davis, E.L.; Baum, T.J. and Hussey, R.S. (2004).** Use of solid-phase sub active hybridization for the identification of parasitism gene candidates from the root-knot nematode, *Meloidogyne incognita*. *Molecular Plant Pathology*, 5: 217-222.
- Kaplan, D.T. and Keen N.T. 1980.** Mechanisms conferring plant incompatibility to nematodes. *Revue Nématol.* 3: 123-134.
- Karajeh, M.R. 2008.** Interaction of the root knot nematode (*Meloidogyne javanica*) and tomato as affected by hydrogen peroxide. *Journal of Plant Protection Research* 48(2): 181-187.
- Lazarovits, G. and Ward, W.B. 1982.** Polyphenoloxidase activity in soybean hypocotyles at sites inoculated with *Phytophthora mega sperma* f.sp. *glycinea*. *Physiol. Plant Pathol.* 21: 227-236.
- Linford, M.B. and Olievera, J.M. (1940).** *Rotylenchulus reniformis*, nov. gen., n. sp. a nematode parasite of roots. *Proc. Helminth. Soc. Wash.* 7: 35-42
- Mannervik, B. and Guthenberg, C. 1981.** Glutathione transferase (Humanplacenta). *Methods in Enzymology*, 77: 231-235.
- Mellersh, D.G.; Foulds, I.V.; Higgins, V.J. and Heath, M. (2002).** H₂O₂ plays different roles in determining penetration failure in three diverse plant-fungal interactions. *New Phytologist*, 170: 501-512.
- Montes, M.J.; Lopez-Brana, I. and Delibes, A. (2004).** Root enzyme activities associated with resistance to *Heterodera avenae* conferred by gene CRE7 in wheat, *Aegilops triuncialis* introgression line. *Journal of Plant Physiology*, 161: 493-495.
- Nakano, Y. and Asada, K. 1981.** Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.*, 22: 867-880.
- Saeed, M.R.M. (2005).** Utilization of some specific materials to stimulate resistance in some host plants against the root-knot nematodes. Ph. D. Thesis. Cairo Univ. Fac. of Agric., 220p.
- Siahpoush, S.; Sahebani, N. and Aminian, H. 2011.** Change in some defense compounds of cucumber treated with *Bacillus cereus* and salicylic acid against *Meloidogyne javanica*. *African J. of Plant Sci.* %(14): 829-834.

- Singh, N.; Ma, L.Q.; Srivastava, M. and Rathinasabapathi, B. 2006.** Metabolic adaptations to arsenic-induced oxidative stress in *Pteris vittata* L. and *Pteris ensiformis* L. *Plant Science*, 170: 274-282.
- Zacheo, G.; Arrigoni-Liso, R.; Beleve-Zacheo, T.; Lamberti, F. and Arrigoni, O. 1983.** Mitochondrial peroxidase and superoxide dismutase activities during infection by *Meloidogyne incognita*. *Nematologica Mediterranea* 10 : 75-80.
- Zacheo, G.; Beleve-Zacheo, T. and Pricolo, G. 1987.** Metabolic changes in enzyme levels in potato roots infested by potato cyst nematode, *Globodera pallid* (Pa3) and *Globodera rostochiensis* (R01). *Nematologica Mediterranea* 15(2):293-302.
- Zacheo, G.; Orlando, C. And Beleve-Zacheo, T. (1993).** Characterization of anionic peroxidase in tomato isolate infected by *Meloidogyne incognita*. *Journal of Nematology*, 25 (2): 249-259.

الملخص العربي

نشاط إنزيمات الأكسدة ومضادات الأكسدة المتعلقة بآليات المقاومة لنيماتودا تعقد الجذور

والنيماتودا الكلوية ونيماتودا الموالح في عوائلها النباتية

عبد المنعم عفيفي*، السيد أبو المعاطي السيد**، نميرأنيس محفوض**، أحمد عبد السلام فرحات**

* قسم الكيمياء الحيوية - كلية الزراعة - جامعة القاهرة

** قسم الحيوان والنيماتولوجيا الزراعية - كلية الزراعة - جامعة القاهرة

أجري هذا البحث بغرض مقارنة التغيرات في نشاط إنزيمات الأكسدة والمضادة للأكسدة وكذلك التغير في المواد المؤكسدة ومضادات الأكسدة في كل من اللوبيا والباذنجان والياسمين والباباؤن نتيجة إصابتها بأي من نيماتودا تعقد الجذور والنيماتودا الكلوية وكذلك في النارج نتيجة إصابته بنيماتودا الموالح. أوضحت النتائج أن رد فعل هذه النباتات للإصابة بأنواع النيماتودا الثلاثة كان تقريباً واحد حيث زاد نشاط الانزيم المؤكسد للدهون وفوق أكسيد الهيدروجين أولاً نتيجة الإصابة مما أدى إلى زيادة المواد المضادة للأكسدة مثل الجلوتاثيون وحمض الأسكوربيك وأيضاً زيادة في نشاط الإنزيمات المضادة للأكسدة مثل سوپر اوكسيد ديسميوتيز، الكاتاليز، اسكوربيك بيروكسيدز والجلوتاثيون-س-ترانسفيريز، غير أن معدل التغير ونسبة الزيادة اختلفت باختلاف نوع النيماتودا ونوع النبات ومستوى العدوى.